

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/235968719>

Albinism in the fishing cat (*Prionailurus viverrinus*) from the Haor Basin of Bangladesh

Article · January 2013

CITATION

1

READS

98

3 authors, including:



[Anthony J. Giordano](#)

Texas Tech University

82 PUBLICATIONS 212 CITATIONS

[SEE PROFILE](#)



[Mohammed Feeroz](#)

Jahangirnagar University

69 PUBLICATIONS 383 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Rarest Cat in the World? Assessing the Conservation Status of the Javan Fishing Cat [View project](#)

ISSN 1027-2992

CAT news

N° 58 | SPRING 2013





CATnews is the newsletter of the Cat Specialist Group, a component of the Species Survival Commission SSC of the International Union for Conservation of Nature (IUCN). It is published twice a year, and is available to members and the Friends of the Cat Group.

**For joining the Friends of the Cat Group please contact
Christine Breitenmoser at ch.breitenmoser@kora.ch**

Original contributions and short notes about wild cats are welcome

**Send contributions and observations to
ch.breitenmoser@kora.ch.**

Guidelines for authors are available at www.catsg.org/catnews

CATnews is produced with financial assistance from Friends of the Cat Group.

Design: barbara surber, werk'sdesign gmbh

Layout: Christine Breitenmoser

Print: Stämpfli Publikationen AG, Bern, Switzerland

ISSN 1027-2992 © IUCN/SSC Cat Specialist Group

The designation of the geographical entities in this publication, and the representation of the material, do not imply the expression of any opinion whatsoever on the part of the IUCN concerning the legal status of any country, territory, or area, or its authorities, or concerning the delimitation of its frontiers or boundaries.



Editors: Christine & Urs Breitenmoser
Co-chairs IUCN/SSC
Cat Specialist Group
KORA, Thunstrasse 31, 3074 Muri,
Switzerland
Tel ++41(31) 951 90 20
Fax ++41(31) 951 90 40
<u.breitenmoser@vetsuisse.unibe.ch>
<ch.breitenmoser@kora.ch>

Associate Editors: Keith Richmond
Brian Bertram
Sultana Bashir
Javier Pereira

Cover Photo: First photographic evidence
of a Pallas's cat in Bhutan. The animal
was captured in Jigme Dorji National
Park on 17.11.2012
Photo: Jigme Dorji National Park

Albinism in fishing cats from the Haor Basin of Bangladesh

The fishing cat *Prionailurus viverrinus* is a small to medium-sized felid that ranges across southern Asia from Pakistan east to Vietnam and south to the island of Java. To date very little is known about the ecology and natural history of this species. Here we document an adult albino fishing cat believed to have been captured in the Hail Haor region of Sylhet Division, Bangladesh and propose that albinism may be well established among fishing cats in the Haor Basin. We are unaware of any other aberrant pelage patterns that have been documented in fishing cats.

The fishing cat is a robust, small to medium-sized felid inhabiting wetland and riparian areas across southern and southeast Asia. Typical pelage for the species consists of brownish-grey to olive-grey fur marked with nearly parallel lines of black, broken elongated spots on the back and sides (Sterndale 1884). Spots vary in shape from rounded to elongate (Francis 2008); the tail is banded or ringed, and white fur is present around the mouth and muzzle. The ears are black and have a conspicuous central white spot on the dorsal surface like its congeneric, the leopard cat *Prionailurus bengalensis* (Roberts 2005). Currently, the fishing cat's ecology and distribution largely remain unknown.

In December 2009 while conducting surveys for fishing cats in northeastern Bangladesh,

we documented the occurrence of an albino male fishing cat (Fig. 1) in a captive wildlife collection in the town of Srimangal. The cat, a male held in captivity for approximately 10 years (captured in August 2001), was captured as an adult at Hail Haor (24°22'N/91°40'E), a large wetland that is currently the focus of our research. The second author examined and photographed the same individual in 2001 a few weeks after its capture.

Khan (2005) reported four albino fishing cats (3 kittens, 1 adult) caught on three occasions within an 18-month period from this same region (2000-2001) suggesting that albinism may be somewhat established in this population. The fate of the three other kittens that Khan (2005) referred to is unclear, as no further information is available. During

our visit, staff managing this captive wildlife collection claimed that other individuals had been observed more recently from the same general region, which suggests anecdotally at least that this trait is still represented in the gene pool.

Considering its time in captivity and its adult status at the time of capture, this albino male was at least 12 years old at the time we observed it. To our knowledge, the Haor region of Sylhet is the only region from which albino fishing cats have been documented. Albinism has been documented in several other felid species (Robinson 1970a, 1976; Nowell & Jackson 1996), particularly in captivity. However, it appears to occur infrequently in wild felid populations. Other somewhat familiar pelage variants in felids include both leucism (Robinson 1969, McBride & Giordano 2010) and melanism (Robinson 1970b, Wibisono & McCarthy 2010, Bashir et al. 2011, Giordano et al. 2012), of which the latter variant has recently been the subject of increasing genetic and field research (Eizirik et al. 2003, Haag et al. 2010, Kawanishi et al. 2010). Although we are not aware of the existence of other fishing cat pelage variants, melanism does occur in leopard cats from the Sundarbans (Kumar Das et al. 2012). As a final thought, it would be of notable scientific interest if albinism does indeed persist in any wild population of felids. Al-



Fig. 1. Albino fishing cat captured in the vicinity of Hail Haor region of Sylhet, Bangladesh (Photo Ali Reza & Anthony J. Giordano).

binism likely has selective disadvantages for predator and prey of numerous taxa, such as causing individuals to be highly conspicuous, and/or being associated with additional phenotypic anomalies. Such characteristics have the potential to lead to or further be associated with overall reduced fitness (DeWoody 2005) and may be more likely to occur in small inbred populations (Bensch et al. 2000). However, few studies have sought to assess these characteristics in the field. It is possible that should albinism persist even infrequently in the Hail Haor fishing cat population, selective agents against the trait do not operate strongly. Fishing cats in the region appear to prey largely on fish and other aquatic prey at night, an ecological context where the importance of pelage coloration might be minimized. Furthermore, fishing cats are currently the largest felid in the region and might not have other natural predators. Finally, it is possible that the population has suffered from some level of inbreeding and is effectively isolated from other nearby fishing cat populations. Though admittedly the above is only speculation at this point, our current and future field work in the region may shed more light on this situation.

References

- Bashir T., Bhattacharya T., Poudyal K., & Sa-thyakumar S. 2011. Notable observations on the melanistic Asiatic golden cat (*Pardofelis temminckii*) of Sikkim, India. *Nebio* 2, 1-4.
- Bensch S., Hansson B., Hasselquist B., & Nielsen B. 2000. Partial albinism in a semi-isolated population of great reed-warblers. *Hereditas* 133, 167-170.
- DeWoody J. A. 2005. Molecular Approaches to the study of parentage, relatedness, and fitness: practical applications for wild animals. *Journal of Wildlife Management* 69, 1400-1418.
- Eizirik E., Yuhki N., Johnson W. E., Menotti-Raymond M., Hannah S. S., & O'Brien S. J. 2003. Molecular genetics and evolution of melanism in the cat family. *Current Biology* 13, 448-453.
- Francis C. M. 2008. A Guide to the Mammals of Southeast Asia. Princeton University Press, Princeton, New Jersey, U.S.A. 392 pp.
- Giordano A. J., de la Cadena E., & Hocking P. 2012. Occurrence of a melanistic pampas cat in the central Peruvian Andes. *Cat News* 57, 14.
- Haag T., Santos A. S., Valdez F. P., Sana D. A., Silveira L., Cullen L., De Angelo C., Morato, R. G., Crawshaw P. G., Salzano F. M., & Eizirik E. 2010. Molecular tracking of jaguar melanism using faecal DNA. *Conservation Genetics* 11, 1239-1242.
- Kawanishi K., Sunquist M. E., Eizirik E., Lynam A. J., Ngoprasert D., Shahruddin W. N. W., Rayan D. M., Sharma D. S. K., & Steinmetz R. 2010. Near fixation of melanism in leopards of the Malay Peninsula. *Journal of Zoology* 282, 201-206.
- Khan M. M. H. 2005. Status and distribution of wild cats in Bangladesh. *Bangladesh Journal of Life Sciences* 17, 69-74.
- Kumar Das S., Kumar Sakar P., Saha R., Vyas P., Danda A. A., & Vattakavan J. 2012. First photographic capture of melanistic leopard cat in Sundarbans. *Cat News* 57, 30-31.
- McBride R. T. & Giordano A. J. 2010. First record of a white ocelot (*Leopardus pardalis*). *The Southwestern Naturalist* 55, 567-568.
- Nowell K. & Jackson P. 1996. Wild Cats, Status Survey and Conservation Action Plan. IUCN/SSC Cat Specialist Group, Gland, Switzerland. 382 pp.
- Roberts T. J. 2005. Field guide to the large and medium-sized mammals of Pakistan. Oxford University Press, Oxford, U.K. 259pp.
- Robinson R. 1969. The white tigers of Rewa and gene homology in the Felidae. *Genetica* 40, 198-200.
- Robinson R. 1970a. Homologous mutants in mammalian coat colour variation. *Symposium of the Zoological Society London* 26, 251-269.
- Robinson R. 1970b. Inheritance of the black form of the leopard (*Panthera pardus*). *Genetica* 41, 190-197.
- Robinson R. 1976. Homologous genetic variation in the Felidae. *Genetica* 46, 1-31.
- Sterndale R. A. 1884. Natural History of the Mammalia of India and Ceylon. Thacker, Spink, and Co., Calcutta, India. 571 pp.
- Wibisono H. T. & McCarthy J. 2010. Melanistic marbled cat from Bukit Barisan Selatan National Park, Sumatra, Indonesia. *Cat News* 52, 9-10.

¹ S.P.E.C.I.E.S., LifeScape International, New York, NY, U.S.A.
*<species@hotmail.com>

² Dept. of Natural Resources Management, Texas Tech University, Lubbock, TX 79409, U.S.A.

³ Division of Biological and Physical Sciences, Delta State University, Cleveland, MS 38733, U.S.A.

⁴ Wildlife Program, Department of Zoology, Jahangirnagar University, Savar, Dhaka 1342, Bangladesh